

Countries/areas reporting vector-borne Zika virus transmission

A. First Zika outbreak since 2015, with no evidence of past circulation, and with ongoing transmission by mosquitoes (44)

High risk of Guillain-Barré syndrome (GBS), microcephaly and other neurological disorders associated with Zika virus

Africa:	Cape Verde
Americas (37):	Aruba, Barbados, Belize, Bolivia, Bonaire, Brazil, Colombia, Costa Rica, Cuba, Curacao, Dominica, Dominican Republic, Ecuador, El Salvador, French Guiana, Guadeloupe, Guatemala, Guyana, Haiti, Honduras, Jamaica, Martinique, Mexico, Nicaragua, Panama, Paraguay, Peru, Puerto Rico, Saint Barthelemy, Saint Lucia, Saint-Martin, St Maarten, St Vincent and the Grenadines, Suriname, Trinidad and Tobago, the US Virgin Islands, Venezuela
Southeast Asia:	Maldives
Western Pacific (5):	American Samoa, Fiji, Marshall Islands, Samoa, Tonga

B. With evidence of Zika transmission before 2015, with or without ongoing transmission; or outbreaks since 2015 reported to be over (17)

Zika transmission at low levels in the past

Africa:	Gabon
Southeast Asia (3):	Bangladesh, Indonesia, Thailand
Western Pacific (12):	Cambodia, Cook Islands*, French Polynesia*, Laos, Malaysia, Micronesia, New Caledonia, Papua New Guinea, Philippines, Solomon Islands, Vanuatu, Vietnam
Americas:	Easter Island (Chile)*

Countries/areas reporting non vector-borne Zika virus transmission

With evidence of person-to-person Zika transmission, other than mosquito-borne (9)

Americas (5):	Argentina, Canada, Chile, Peru, United States
Europe (3):	France, Italy, Portugal
Western Pacific:	New Zealand

**Outbreaks since 2015 reported over*