

## Appendix

Incubation period (period from infection to appearance of illness) of measles ranges from seven days to up to 21 days. The general public who has visited the public places during the period listed below should observe if they develop relevant symptoms, such as fever, skin rash, cough, runny nose and red eyes in seven to 21 days (medical surveillance period). If symptoms arise, they should wear surgical masks, stop going to work or school and avoid going to crowded places. They should also avoid contact with non-immune persons, especially persons with weakened immunity, pregnant women and children aged below one. Those suspected to have infected are advised to seek medical attention as early as possible. If they need to visit any health care facilities during the period of medical surveillance, they should also report whether they have symptoms and prior measles exposure history to the healthcare workers so that appropriate infection control measures can be implemented at the concerned healthcare facilities to prevent any potential spread.

List of public places the measles patient visited during the period of communicability:

	Date	Time	Place	End of medical surveillance for contacts
Female / aged 25	June 10	09:41-10:00	Human Health Medical Centre Shop 2C, G/F, Bo Fung Gardens, 1090 King's Road, Quarry Bay	July 1
	June 10	16:15-17:00	Quality HealthCare Medical Centre Suite 2405 - 06, 24/F World Trade Centre, 280 Gloucester Road, Causeway Bay	July 1
	June 10	20:38-00:00	Accident and Emergency Department at United Christian Hospital	July 1
	June 11	00:01-07:05	Accident and Emergency Department at United Christian Hospital	July 2