Appendix

Incubation period (period from infection to appearance of illness) of measles ranges from seven days to up to 21 days. The general public who has visited the public places during the period listed below should observe if they develop relevant symptoms, such as fever, skin rash, cough, runny nose and red eyes in seven to 21 days (medical surveillance period). If symptoms arise, they should wear surgical masks, stop going to work or school and avoid going to crowded places. They should also avoid contact with non-immune persons, especially persons with weakened immunity, pregnant women and children aged below one. Those suspected to have infected are advised to seek medical attention as early as possible. If they need to visit any health care facilities during the period of medical surveillance, they should also report whether they have symptoms and prior measles exposure history to the healthcare workers so that appropriate infection control measures can be implemented at the concerned healthcare facilities to prevent any potential spread.

List of public places the measles patient visited during the period of communicability:

	Date	Time	Place	End of medical
				surveillance for
				contacts
Male /	Oct 1	06:20-08:00	Lok Ma Chau Spur Line Control	Oct 22
aged 30			Point	
		08:00-13:30	Lo Wu Control point	
		13:30-14:30	Lok Ma Chau Spur Line Control	
			Point	
	Oct 2	14:30-15:30	Lok Ma Chau Spur Line Control	Oct 23
			Point	
		15:30-17:30	Lo Wu Control point	
		17:30-22:45	Lok Ma Chau Spur Line Control	
			Point	
	Oct 4	06:20-08:00	Lok Ma Chau Spur Line Control	Oct 25
			Point	
		08:00-13:30	Lo Wu Control point	
		13:30-14:30	Lok Ma Chau Spur Line Control	
			Point	
	Oct 5	09:00-09:30	Yau Ma Tei Jockey Club General Out-patient Clinic	Oct 26