

**Annual concentration averages of the pollutants in the monitoring network from  
2006 to 2019**

Year	Annual concentration averages of pollutants					
	Sulphur dioxide ( $\mu\text{g}/\text{m}^3$ )	Nitrogen dioxide ( $\mu\text{g}/\text{m}^3$ )	Ozone ( $\mu\text{g}/\text{m}^3$ )	Respirable suspended particulates ( $\mu\text{g}/\text{m}^3$ )	Fine suspended particulates ( $\mu\text{g}/\text{m}^3$ )	Carbon monoxide ( $\text{mg}/\text{m}^3$ )
2006	43	42	44	67	-	-
2007	44	41	46	72	-	-
2008	36	40	46	65	-	-
2009	26	38	51	64	-	-
2010	23	39	49	59	-	-
2011	21	37	53	59	-	-
2012	17	35	49	52	-	-
2013	17	37	49	59	-	-
2014	14	34	52	50	-	-
2015	12	30	47	44	29	0.730
2016	11	32	44	41	26	0.728
2017	10	31	52	45	28	0.665
2018	9	29	53	42	25	0.611
2019	7	30	60	42	25	0.700

Note: The concentrations of gaseous pollutants are calculated at a reference temperature of 298.15K and a pressure of 101.325 kPa and the concentrations of PM<sub>10</sub> and PM<sub>2.5</sub> are measured at real-time temperature and atmospheric pressure